

THE NATURAL SOLUTION THAT BOOSTS PROTECTION AND SECURES YIELD



FLEXIBILITY

Compatible in mixtures*
Pre Harvest Interval (PHI) = 0 day

*Please contact us for more information

SECURITY

Anti-fungal protection Safe for the user and the environment

SUSTAINABILITY

No residue Biodegradable Additional mode of action

Recommended between 25 to 200 gal/acre.

SPRAY VOLUME

Initial spray should be made before or at the first sign of disease and then every 7 to 14 days during the infection period.

TIMING / FREQUENCY

CUCUMBER DOWNY MILDEW TRIAL YIELD AT HARVEST - 2023

Small plot trials (Randomized Complete Block, 4 replications)

Variety: Diamondback

Contract research organization: Glades Crop Care

Stone fruits - such as peaches, plums, nectarines,

cherries and hybrids, etc.

Location: Jupiter (FL), USA


MUSKMELON DOWNY MILDEW TRIAL **DISEASE INCIDENCE AT HARVEST - 2023**


Small plot trials (Randomized Complete Block, 4 replications)

Variety: Hales Best

Contract research organization: Aq-Master Research

Location: Richmond (TX), USA







Liquid formulation, easy to use No storage constraints Shelf life: 3 years



6 to 20 fl.oz./acre

